

Program:

1. C^* -algebras: definition and examples.
2. Unitalization of a C^* -algebra.
3. Spectrum of an element of a C^* -algebra, its properties.
4. Commutative C^* -algebras. The space of maximal ideals. The Gelfand transform.
5. Gelfand's theorem about commutative C^* -algebras.
6. The Stone-Weierstrass theorem.
7. C^* -algebra generated by a normal element. The functional calculus for normal operators.
8. Positive elements, its properties.
9. Approximate units, their existence.
10. Ideals, factor-algebras, hereditary subalgebras.
11. Automatic continuity of $*$ -homomorphisms.
12. Von Neumann algebras. Bicommutant theorem.
13. Topologically irreducible representations.
14. Positive functionals, states.
15. GNS-construction.
16. Realization of C^* -algebras as operator algebras (Gelfand-Naimark theorem).
17. Jordan decomposition.
18. Finite dimensional linear topological spaces. Uniqueness of Hausdorff topology.
19. Finite-dimensional C^* -algebras, their unitality and structure.
20. Non-degenerate representations.
21. The algebra of compact operators and its properties.
22. AF-algebras, a description of homomorphisms of finite-dimensional algebras, Bratteli diagrams.

Additional list of problems (to formulated at lectures)

1. Let A be a C^* -algebra, $a \in A$, $p, q \in A$ — orthogonal projections (i.e. self-adjoint idempotents with $pq = 0$). Show that if a is positive and $pap = 0$, then $paq = 0$.
2. Let A be a C^* -algebra, $a \in A$. Let us denote by aAa the set of all elements of the form aba , where $b \in A$, and by \overline{aAa} the closure of this set. A C^* -subalgebra $B \subset A$ is *hereditary* if the conditions $0 \leq a \leq b$ and $b \in B$ imply that $a \in B$.
 - (a) Check that \overline{aAa} is a C^* -subalgebra for any $a \in A$.
 - (b) Let $p \in A$ be a projection. Verify that pAp is closed.
 - (c) Show that pAp is hereditary for any projector p .
 - (d) Show that \overline{aAa} is hereditary for any positive $a \in A$.
3. Let $X \subset \mathbb{R}$ be the set of points $1, 1/2, 1/3, \dots$ and 0 . Let $C(X, M_2)$ be the set of all continuous functions on X with values in the matrix algebra M_2 . Let $B_1 = \{f \in C(X, M_2) : f(0) \text{ is diagonal}\}$, $B_2 = \{f \in C(X, M_2) : f(0) \text{ has the form } \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}\}$.
 - (a) Show that $C(X, M_2)$, B_1 , B_2 are C^* -algebras.
 - (b) Find all (two-sided, closed) ideals in $C(X)$, $C(X, M_2)$, B_1 , B_2 .
4. Let A be a C^* -algebra, $J \subset A$ be an ideal, $a \in A$ is a self-adjoint element. Show that there exists an element $j \in J$ such that $\|[a]\| = \|a - j\|$, where $[a] \in A/J$ is the class $a + J$ of element a . Hint: decompose $a - \|[a]\| \cdot 1 = a_+ - a_-$ with positive a_+ , a_- and show that $a_+ \in J$.
5. Let A be a C^* -algebra, $a \in A$ be a self-adjoint element. Show that if the spectrum $\sigma(a)$ is an infinite set, then A is infinite-dimensional.
6. Describe the GNS construction for the C^* -algebra $C[0, 1]$ and for a positive linear functional φ
 - (a) $\varphi(f) = f(0)$,
 - (b) $\varphi(f) = \frac{1}{2}(f(0) + f(1))$,
 - (c) $\varphi(f) = \int_0^1 f(x) dx$,
 where $f \in C[0, 1]$.
7. Describe the GNS construction for the C^* -algebra M_n of complex $n \times n$ -matrices and for a positive linear functional φ
 - (a) $\varphi(A) = a_{11}$,
 - (b) $\varphi(A) = \text{tr}(A)$,
 where $A = (a_{ij})_{i,j=1}^n \in M_n$.

8. Let π, σ be representations of a C^* -algebra A on the Hilbert spaces H_π and H_σ , and let a partial isometry $U : H_\pi \rightarrow H_\sigma$ satisfy the equality $\sigma(a)U = U\pi(a)$ for any $a \in A$. Show that the image (resp. orthogonal complement to the kernel) of U is an invariant subspace for $\sigma(A)$ (resp. for $\pi(A)$). (U is a partial isometry if U^*U and UU^* are projections)
9. (a) Let $M_n(A)$ be the set of all $n \times n$ -matrices with coefficients from a C^* -algebra A . Show that on $M_n(A)$ there exists a C^* -norm.
- (b) Let A be a C^* -algebra with norm $\|\cdot\|$, and let $\|\cdot\|'$ be another norm on A , equivalent to the first norm. Show that if $\|\cdot\|'$ is a C^* -norm, then these norms coincide. Deduce from this the uniqueness of C^* -norm on $M_n(A)$.
10. Let φ be a state on a C^* -algebra A . Suppose that for some self-adjoint element $a \in A$ one has the equality $\varphi(a^2) = \varphi(a)^2$. Show that it follows from this that $\varphi(ab) = \varphi(ba) = \varphi(a)\varphi(b)$ for any $b \in A$.
11. Let $A = c$ be the C^* -algebra of all convergent sequences of complex numbers, $c = \{(a_n)_{n \in \mathbb{N}} : a_n \in \mathbb{C}; \lim_{n \rightarrow \infty} a_n \text{ exists}\}$. Let us consider it as a C^* -subalgebra of the algebra $\mathbb{B}(l_2)$ of bounded operators in the Hilbert space l_2 of square-integrable sequences. Find the first and second commutant, A' and A'' , and (independently) the weak closure of A in $\mathbb{B}(l_2)$.
12. (a) Show that the weak topology is strictly weaker than the strong topology.
- (b) Let $P \subset \mathbb{B}(H)$ be the set of all (self-adjoint) projections on a Hilbert space. Show that if $p_\lambda \rightarrow p$ weakly converges, where $p_\lambda \in P$ and $p \in P$, then $p_\lambda \rightarrow p$ strongly converges.
- (c) Show that the strong limit of a sequence of (self-adjoint) projections is a projection.
- (d) Find an example of a weakly convergent net $p_\lambda \rightarrow p$ with $p_\lambda \in P$ and $p \notin P$.
13. Let $H_n \subset H$ be the subspace of a Hilbert space H generated by the first n vectors of an orthonormal basis. In the set of all sequences (m_1, m_2, \dots) , where $m_k \in \mathbb{B}(H_n) \subset \mathbb{B}(H)$, consider the subset A of all sequences such that
- $\sup_k \|m_k\| < \infty$;
 - the sequences (m_1, m_2, \dots) and (m_1^*, m_2^*, \dots) are convergent in the strong topology.
- Show that A is a C^* -algebra and that the mapping $(m_1, m_2, \dots) \mapsto s\text{-}\lim_{k \rightarrow \infty} m_k \in \mathbb{B}(H)$ is a surjective $*$ -homomorphism of $A \rightarrow \mathbb{B}(H)$.
14. Let A be a commutative C^* -algebra and let π be its irreducible representation on a Hilbert space H . Show that $\dim H = 1$
15. Consider $C[0, 1]$ as a C^* -subalgebra in $\mathbb{B}(H)$, where $H = L^2([0, 1])$ (continuous functions act on H by multiplication).

- (a) Check that $C[0, 1] \cap \mathbb{K}(H) = 0$;
- (b) Let φ be a linear functional on $C[0, 1]$ defined by the equality $\varphi(f) = f(0)$, $f \in C[0, 1]$. Find a sequence of $\{e_n\}_{n \in \mathbb{N}}$ vectors of unit length weakly converging to zero in H such that $\varphi(f) = \lim_{n \rightarrow \infty} \langle fe_n, e_n \rangle$ for any function $f \in C[0, 1]$.
16. Operators a, b in a Hilbert space H are called *compalent* if there exists a unitary operator $u \in \mathbb{B}(H)$ such that $u^*au - b \in \mathbb{K}(H)$. Show that if self-adjoint operators a, b are compalent then their essential spectra coincide.
17. Show that any AF C^* -algebra without unity has an approximative unity consisting of an increasing sequence of projections.
18. (a) Show that $C[0, 1]$ is not an AF-algebra.
- (b) Construct an injective $*$ -homomorphism $C[0, 1]$ into the AF-algebra $C(K)$ of continuous functions on the Cantor set K . Hint: construct a function f on K that takes all rational values from $[0, 1]$ and show that $C^*(f)$ is isometrically $*$ -isomorphic $C(\text{Sp}(f)) = C[0, 1]$.
19. Let $A_n = M_{2^n}(\mathbb{C}) \oplus M_{2^n}(\mathbb{C})$, and let the embedding $\alpha_n : A_n \rightarrow A_{n+1}$ be given by the formula $\alpha_n : \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix} \mapsto \begin{pmatrix} a_1 & 0 & | & 0 & 0 \\ 0 & a_1 & | & 0 & 0 \\ \hline 0 & 0 & | & a_1 & 0 \\ 0 & 0 & | & 0 & a_2 \end{pmatrix}$, where $a_1, a_2 \in M_{2^n}(\mathbb{C})$.
- (a) Find the Bratteli diagram for the AF algebra $A = \overline{\bigcup_{n=1}^{\infty} A_n}$;
- (b) Find whether A is unital.