

Problem 1. Give an example of an involutive Banach algebra that is not a C^* -algebra.

Problem 2. Show that a left unit element is also a right one, which has $1^* = 1$; that the identity element is unique and that $\|1\| = 1$. It is called the *unit of algebra*.

Problem 3. Verify that the algebra $C(X)$ formed by all continuous complex-valued functions on a compact space X and the algebra $C_0(X)$ of all continuous complex-valued functions on a locally compact space X tending to 0 at infinity (that is, $f : X \rightarrow \mathbb{C}$ such that for any $\varepsilon > 0$ there exists a compact $K \subseteq X$ such that $\sup\{|f(x)| \mid x \in K\} < \varepsilon$) are commutative C^* -algebras if the *supremum-norm*: $\|f\| = \sup_{x \in X} |f(x)|$, is taken as the norm and the pointwise multiplication is taken as the multiplication. Moreover, the algebra $C(X)$ is unital.

Problem 4. Verify that the algebra $\mathbb{B}(H)$ of all bounded operators acting on a Hilbert space H is a C^* -algebra with identity. Here as a norm we take the *operator norm* $\|a\| = \sup_{h \in H, \|h\| \leq 1} \|a(h)\|$, and the multiplication is the composition of operators.

Problem 5. Show that “sum” norm turns A^+ into an involutive Banach algebra, but not into a C^* -algebra.

Problem 6. Prove that A is an ideal in A^+ .

Problem 7. Prove that if $a_0 \in A$ is invertible and $\|a - a_0\| < \frac{1}{\|a_0^{-1}\|}$, then a is also invertible, and $a^{-1} = \sum_{n=0}^{\infty} [a_0^{-1}(a_0 - a)]^n a_0^{-1}$.

Problem 8. Show that the quasi-spectrum always contains zero.

Problem 9. Let a and b be commuting elements of a Banach algebra. Then the product ab is invertible if and only if each of the elements a and b are invertible.

Problem 10. Show that $r(a) \leq \|a\|$ for any $a \in A$.

Problem 11. Prove that M_A is a locally compact Hausdorff space, and M_{A^+} is its one-point compactification.

Problem 12. Check that if $\varphi : A \rightarrow \mathbb{C}$ is a (non-zero) multiplicative functional on A , then the formula $\tilde{\varphi}((a, \lambda)) = \varphi(a) + \lambda$ defines a unique extension of φ to a multiplicative functional on A^+ .

Problem 13. Any ideal I of a commutative unital Banach algebra is contained in some maximum ideal.

Problem 14. Prove the Gelfand theorem for a non-unital algebra.

Problem 15. If a is an invertible element, then the algebra $C^*(a)$ is unital. In this case $C^*(a) = C^*(1, a)$.

Problem 16. Prove that $f(\text{Sp}(a)) = \text{Sp}(f(a))$ approximating f by polynomials, and correctly stating what it means that the image is continuous under a uniform approximation, and using the isometricity of the Gelfand transform.

Problem 17. Show that if $a \geq 0$ and $0 \geq a$, then $a = 0$; and also that $-\|a\|1 \leq a \leq \|a\|1$ for every self-adjoint a .

Problem 18. If $0 \leq a \leq b$, then $\|a\| \leq \|b\|$.

Problem 19. Prove that an algebra with countable approximate unit does not have to be separable.

Problem 20. Prove that a positive element of an arbitrary C^* -subalgebra is a positive element of the entire algebra.

Problem 21. Let $\varphi : A \rightarrow B$ be a $*$ -homomorphism of non-unital algebras. Prove that there is a unique unital $*$ -homomorphism $\varphi^+ : A^+ \rightarrow B^+$, extending φ .

Problem 22. Let $\varphi : A \rightarrow B$ be a $*$ -homomorphism of algebras, with A non-unital, and B unital. Prove that there is a unique unital $*$ -homomorphism $\varphi^{(+)} : A^+ \rightarrow B$, extending φ .

Problem 23. Verify that

- If S is self-adjoint, then so is S' .
- The commutant of any set is a unital algebra.
- The commutant of any set is weakly closed.
- Thus, S' is the von Neumann algebra for any self-adjoint set S .
- If $S_1 \subset S_2$, then $S'_1 \supset S'_2$.
- Always $S \subset S''$.
- Therefore $S' = S'''$, $S'' = S''''$, etc.

Problem 24. Prove that if a self-adjoint element a in a unital C^* -algebra has $\text{Sp}(a) = \{0, 1\}$, then a is a nonscalar idempotent.

Problem 25. Let u_λ , $\lambda \in \Lambda$, be some approximate unit in a unital algebra. Prove that $1 = \lim_{\lambda \in \Lambda} u_\lambda$.

Problem 26. There is a natural bijection between self-adjoint linear functionals on A and (real) linear functionals on A_{sa} .

Problem 27. If A is unital, then a representation π is non-degenerate if and only if $\pi(1) = 1$.

Problem 28. Prove that the matrix algebra M_n is simple for any n

Problem 29. Prove that the image of the matrix algebra M_n under a $*$ -homomorphism is either a zero algebra or an algebra isomorphic to M_n .

Problem 30. Draw Bratteli diagram (for some defining sequence) of the following AF-algebra: the algebra of compact operators $\mathbb{K}(H)$.

Problem 31. Draw Bratteli diagram (for some defining sequence) of the following AF-algebra: the unitization $\mathbb{K}(H)^+$ of the algebra of compact operators $\mathbb{K}(H)$.

Problem 32. Draw Bratteli diagram (for some defining sequence) of the following AF-algebra: the closure of the union of $A_p = M_{2^p}$, with embeddings $A_p \subset A_{p+1}$ of multiplicity 2 according to the formula $a \mapsto \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ (CAR algebra).

Problem 33. Draw Bratteli diagram (for some defining sequence) of the following AF-algebra: $C(K)$, where K is the Cantor set obtained from $[0, 1]$ by successive removing the middle third of the corresponding intervals. If K_p is a set, obtained at the p th step of this process, then A_p is an algebra of continuous functions constant on intervals of K_p .

Problem 34. Draw Bratteli diagram (for some defining sequence) of the following AF-algebra: $C(X)$, where $X := \{0\} \cup \{\frac{1}{n} : n \in \mathbb{N}\}$, and A_k consists of all functions constant on $[0, 1/2^k]$.

Problem 35. Let $\pi : A \rightarrow \mathbb{B}(H)$ be a degenerate representation. Let us denote by H_0 the invariant subspace $H_0 := \{\xi \in H : \pi(a)(\xi) = 0 \text{ for any } a \in A\}$. Prove that π induces a representation $\pi' : A \rightarrow \mathbb{B}(H/H_0)$, and if π was a faithful representation (an injective homomorphism), then so is π' .

Problem 36. Let $\pi : A \rightarrow \mathbb{B}(H)$ be a degenerate representation. Let us denote by H_0 the invariant subspace $H_0 := \{\xi \in H : \pi(a)(\xi) = 0 \text{ for any } a \in A\}$. Prove that π induces a representation $\pi' : A \rightarrow \mathbb{B}(H/H_0)$, and if π was a faithful representation (an injective homomorphism), then so is π' .

Problem 37. Check that $RM(A) = (LM(A))^*$ and that $M(A) = LM(A) \cap RM(A)$, so that $M(A)$ is symmetric with respect to the involution.

Problem 38. Prove that a representation $\pi : A \rightarrow \mathbb{B}(H)$ is non-degenerate if and only if for some approximate unit u_λ of the algebra A the following condition is satisfied: for any vector $\xi \in H$ there is a λ such that $u_\lambda(\xi) \neq 0$.

Problem 39. Construct an example of an operator on Hilbert C^* -module that does not admit an adjoint.

Problem 40. Prove that $\|x\| = \sup_{y \in B_1(M)} |\langle x, y \rangle|$, where $B_1(M) \subset M$ is the unit ball.

Problem 41. Prove that $\theta_{x,y} \in \mathbb{B}_A^\star(M)$ by providing an explicit formula for the adjoint.

Problem 42. Prove that $\mathbb{K}_A(M)$ is an ideal in $\mathbb{B}_A^\star(M)$.

Problem 43. Prove that $\mathbb{K}_A(A) = A$. Note that if A is non-unital, then $\mathbb{K}_A(A) = A \neq \mathbb{B}_A^\star(A) = DC(A)$ (algebra of double centralizers).

Problem 44. Let A be a C^* -algebra, $a \in A$, $p, q \in A$ — orthogonal projections (i.e. self-adjoint idempotents with $pq = 0$). Show that if a is positive and $pap = 0$, then $paq = 0$.

Problem 45. Let A be a C^* -algebra, $a \in A$. Let us denote by aAa the set of all elements of the form aba , where $b \in A$, and by \overline{aAa} the closure of this set. A C^* -subalgebra $B \subset A$ is *hereditary* if the conditions $0 \leq a \leq b$ and $b \in B$ imply that $a \in B$.

- (1) Check that \overline{aAa} is a C^* -subalgebra for any $a \in A$.
- (2) Let $p \in A$ be a projection. Verify that pAp is closed.
- (3) Show that \overline{pAp} is hereditary for any projector p .
- (4) Show that \overline{aAa} is hereditary for any positive $a \in A$.

Problem 46. Let $X \subset \mathbb{R}$ be the set of points $1, 1/2, 1/3, \dots$ and 0 . Let $C(X, M_2)$ be the set of all continuous functions on X with values in the matrix algebra M_2 . Let $B_1 = \{f \in C(X, M_2) : f(0) \text{ is diagonal}\}$, $B_2 = \{f \in C(X, M_2) : f(0) \text{ has the form } \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}\}$.

- (1) Show that $C(X, M_2)$, B_1 , B_2 are C^* -algebras.
- (2) Find all (two-sided, closed) ideals in $C(X)$, $C(X, M_2)$, B_1 , B_2 .

Problem 47. Let A be a C^* -algebra, $J \subset A$ be an ideal, $a \in A$ is a self-adjoint element. Show that there exists an element $j \in J$ such that $\|[a]\| = \|a - j\|$, where $[a] \in A/J$ is the class $a + J$ of element a . Hint: decompose $a - \|[a]\| \cdot 1 = a_+ - a_-$ with positive a_+ , a_- and show that $a_+ \in J$.

Problem 48. Let A be a C^* -algebra, $a \in A$ be a self-adjoint element. Show that if the spectrum $\sigma(a)$ is an infinite set, then A is infinite-dimensional.

Problem 49. Describe the GNS construction for the C^* -algebra $C[0, 1]$ and for a positive linear functional φ

- (1) $\varphi(f) = f(0)$,
- (2) $\varphi(f) = \frac{1}{2}(f(0) + f(1))$,
- (3) $\varphi(f) = \int_0^1 f(x) dx$,

where $f \in C[0, 1]$.

Problem 50. Describe the GNS construction for the C^* -algebra M_n of complex $n \times n$ -matrices and for a positive linear functional φ

- (1) $\varphi(A) = a_{11}$,
- (2) $\varphi(A) = \text{tr}(A)$,

where $A = (a_{ij})_{i,j=1}^n \in M_n$.

Problem 51. Let π, σ be representations of a C^* -algebra A on the Hilbert spaces H_π and H_σ , and let a partial isometry $U : H_\pi \rightarrow H_\sigma$ satisfy the equality $\sigma(a)U = U\pi(a)$ for any $a \in A$. Show that the image (resp. orthogonal complement to the kernel) of U is an invariant subspace for $\sigma(A)$ (resp. for $\pi(A)$). (U is a partial isometry if U^*U and UU^* are projections)

Problem 52. (a) Let $M_n(A)$ be the set of all $n \times n$ -matrices with coefficients from a C^* -algebra A . Show that on $M_n(A)$ there exists a C^* -norm.

(b) Let A be a C^* -algebra with norm $\|\cdot\|$, and let $\|\cdot\|'$ be another norm on A , equivalent to the first norm. Show that if $\|\cdot\|'$ is a C^* -norm, then these norms coincide. Deduce from this the uniqueness of C^* -norm on $M_n(A)$.

Problem 53. Let φ be a state on a C^* -algebra A . Suppose that for some self-adjoint element $a \in A$ one has the equality $\varphi(a^2) = \varphi(a)^2$. Show that it follows from this that $\varphi(ab) = \varphi(ba) = \varphi(a)\varphi(b)$ for any $b \in A$.

Problem 54. Let $A = c$ be the C^* -algebra of all convergent sequences of complex numbers, $c = \{(a_n)_{n \in \mathbb{N}} : a_n \in \mathbb{C}; \lim_{n \rightarrow \infty} a_n \text{ exists}\}$. Let us consider it as a C^* -subalgebra of the algebra $\mathbb{B}(l_2)$ of bounded operators in the Hilbert space l_2 of square-integrable sequences. Find the first and second commutant, A' and A'' , and (independently) the weak closure of A in $\mathbb{B}(l_2)$.

- Problem 55.** (a) Show that the weak topology is strictly weaker than the strong topology.
- (b) Let $P \subset \mathbb{B}(H)$ be the set of all (self-adjoint) projections on a Hilbert space. Show that if $p_\lambda \rightarrow p$ weakly converges, where $p_\lambda \in P$ and $p \in P$, then $p_\lambda \rightarrow p$ strongly converges.
- (c) Show that the strong limit of a sequence of (self-adjoint) projections is a projection.
- (d) Find an example of a weakly convergent net $p_\lambda \rightarrow p$ with $p_\lambda \in P$ and $p \notin P$.

Problem 56. Let $H_n \subset H$ be the subspace of a Hilbert space H generated by the first n vectors of an orthonormal basis. In the set of all sequences (m_1, m_2, \dots) , where $m_k \in \mathbb{B}(H_n) \subset \mathbb{B}(H)$, consider the subset A of all sequences such that

- $\sup_k \|m_k\| < \infty$;
- the sequences (m_1, m_2, \dots) and (m_1^*, m_2^*, \dots) are convergent in the strong topology.

Show that A is a C^* -algebra and that the mapping $(m_1, m_2, \dots) \mapsto s\text{-}\lim_{k \rightarrow \infty} m_k \in \mathbb{B}(H)$ is a surjective $*$ -homomorphism of $A \rightarrow \mathbb{B}(H)$.

Problem 57. Let A be a commutative C^* -algebra and let π be its irreducible representation on a Hilbert space H . Show that $\dim H = 1$

Problem 58. Let $\{e_i\}_{i=1}^\infty$ be an orthonormal basis of a Hilbert space H , and let the operators a, b be given by the equalities $ae_i = e_{2i}$; $be_i = e_{2i-1}$. Let $E = C^*(a, b) \subset \mathbb{B}(H)$ be the C^* -algebra generated by a and b .

(a) Check the boundedness of a and b and prove the equalities $a^*a = b^*b = 1$, $aa^* + bb^* = 1$.

(b) Prove that E is not isomorphic to the complete group C^* -algebra $C^*(G)$ for any group G .

Problem 59. Consider $C[0, 1]$ as a C^* -subalgebra in $\mathbb{B}(H)$, where $H = L^2([0, 1])$ (continuous functions act on H by multiplication).

- (a) Check that $C[0, 1] \cap \mathbb{K}(H) = 0$;
- (b) Let φ be a linear functional on $C[0, 1]$ defined by the equality $\varphi(f) = f(0)$, $f \in C[0, 1]$. Find a sequence of $\{e_n\}_{n \in \mathbb{N}}$ vectors of unit length weakly converging to zero in H such that $\varphi(f) = \lim_{n \rightarrow \infty} \langle fe_n, e_n \rangle$ for any function $f \in C[0, 1]$.

Problem 60. Operators a, b in a Hilbert space H are called *compalent* if there exists a unitary operator $u \in \mathbb{B}(H)$ such that $u^*au - b \in \mathbb{K}(H)$. Show that if self-adjoint operators a, b are compalent then their essential spectra coincide.

Problem 61. Show that any AF C^* -algebra without unity has an approximative unity consisting of an increasing sequence of projections.

Problem 62. (1) Show that $C[0, 1]$ is not an AF-algebra.

- (2) Construct an injective $*$ -homomorphism $C[0, 1]$ into the AF-algebra $C(K)$ of continuous functions on the Cantor set K . Hint: construct a function f on K that takes all rational values from $[0, 1]$ and show that $C^*(f)$ is isometrically $*$ -isomorphic $C(\text{Sp}(f)) = C[0, 1]$.

Problem 63. Let $A_n = M_{2^n}(\mathbb{C}) \oplus M_{2^n}(\mathbb{C})$, and let the embedding $\alpha_n : A_n \rightarrow A_{n+1}$ be

given by the formula $\alpha_n : \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix} \mapsto \begin{pmatrix} a_1 & 0 & | & 0 & 0 \\ 0 & a_1 & | & 0 & 0 \\ \hline 0 & 0 & | & a_1 & 0 \\ 0 & 0 & | & 0 & a_2 \end{pmatrix}$, where $a_1, a_2 \in M_{2^n}(\mathbb{C})$.

- (1) Find the Bratteli diagram for the AF algebra $A = \overline{\bigcup_{n=1}^\infty A_n}$;
- (2) Find whether A is unital.

Problem 64. Find $M(A)$, where $A = \{f \in C([0, 1]; M_2) : f(0)_{11} = f(0)_{12} = f(0)_{21} = 0; f(1) = 0\}$ (here M_2 is the algebra of two-dimensional matrices).